

## INFLUENCE OF AROMATIC SUBSTITUENTS ON THE SYNTHESIS OF SCHIFF BASES DERIVED FROM *TRANS*-(*R,R*)- DIAMINOCYCLOHEXANE: A SPECTROPHOTOMETRIC AND DFT B3LYP STUDY

Lili Dahiana Becerra, Carlos Coy-Barrera\*, Diego Quiroga

*<sup>a</sup>Bioorganic Chemistry Laboratory, Faculty of Basic and Applied Sciences, Nueva Granada Campus, Nueva Granada Military University, Cajicá 250247, Colombia*

*\*e-mail: carlos.coy@unimilitar.edu.co*

**Abstract.** In this work, the synthesis of Schiff bases derived from *trans*-(*R,R*)-diaminocyclohexane by microwave irradiation (MW) is presented. The reaction yields varied between 31% and 69%, being influenced by the electronic nature of the substituents (H, Cl, Br, NO<sub>2</sub>, MeO, *t*-BuO, BnO, and 4-(4-Me)PhO) and the reaction temperature. The spectrophotometric properties of the products were investigated by UV-Vis spectrophotometry, revealing bathochromic and *hypso*-chromic effects attributable to the different substituent groups. These effects were interpreted by DFT calculations with the B3LYP functional at the 6-311G(d,p) level. The results suggest that the electronic properties of the substituents in the para position have a significant impact on the spectroscopic characteristics of the Schiff bases. The synthesized Schiff bases exhibit great potential for applications in areas such as optical sensors and functional materials, as the substituents can precisely modulate their spectrophotometric properties. This opens up new opportunities for designing compounds with tunable properties for various technological and scientific applications.

**Keywords:** schiff base, MW irradiation, substituent effect, spectrophotometric property, DFT calculation.